

## **A Green Morph of the Migratory Locust, *Locusta migratoria* L. (Orthoptera: Acrididae) that Occurred After Inbreeding**

Authors: Tanaka, Seiji, and Nishide, Yudai

Source: Journal of Orthoptera Research, 21(2) : 175-177

Published By: Orthopterists' Society

URL: <https://doi.org/10.1665/034.021.0205>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# A green morph of the migratory locust, *Locusta migratoria* L. (Orthoptera: Acrididae) that occurred after inbreeding

SEIJI TANAKA AND YUDAI NISHIDE

Locust Research Laboratory, National Institute of Agro-biological Sciences at Ohwashi, Tsukuba, Ibaraki 305-8634, Japan.  
Email: stanaka@affrc.go.jp

## Abstract

Green-colored hatchlings appeared in the 2<sup>nd</sup> generation of inbred family lines in *Locusta migratoria* L. Female adults collected in Tsukuba, Japan were kept individually and eggs collected from them. Locusts of the first generation from each female were reared in a group and eggs of the 2<sup>nd</sup> generation obtained from them. Green and normal-colored (fawn) hatchlings appeared together from the same egg pods, originating from 2 of the 34 family lines established. When they were reared together in the same cage, the green morph suffered from a high rate of mortality. A few green-morph female individuals attained the adult stage with light body coloration, but died without producing any eggs. Although no information is available about their genetic background, it seems likely these green morphs represent a recessive, semilethal trait that appeared after inbreeding of wild-caught female adults. Photographs of hatchlings, nymphs and adults are presented.

## Key words

body color, eye color, inbreeding, *Locusta migratoria*

## Introduction

The migratory locust, *Locusta migratoria* L., is widely distributed in the Old World (Uvarov 1966, 1977). This locust feeds on grass and often causes serious damage to agricultural crops. It exhibits remarkable phenotypic plasticity called density-dependent polyphenism and has attracted much attention from many researchers (Pener 1991, Pener & Yerushalmi 1998, Pener & Simpson 2009). It occurs usually in the solitary phase at low population densities and does not cause any economic problem. Nymphs display various cryptic body colors. As population densities increase under favorable conditions, nymphs start aggregating and develop black patterns with a dirty orange color. Hatchling body color is also different between the solitary and gregarious phases and depends on parental crowding conditions. Eggs laid by solitary females give rise to pale-colored hatchlings, whereas those laid by gregarious females produce dark-colored hatchlings (Hunter-Jones 1958). During a study on phase polyphenism in *L. migratoria*, we obtained hatchlings with unusual body coloration in inbred colonies and reared them in the laboratory. This paper describes their body coloration and development.

## Materials and methods

Approximately 40 adult females were collected in grassland in Tsukuba, Ibaraki, Japan in October, 2010. They were apparently in the solitary phase, at a population density < 1/100m<sup>2</sup>. They were individually reared in small screen cages (28 × 15 × 28 cm) at

30 °C with a LD 12:12 hour photoperiod, by a method described previously (Hakomori & Tanaka 1992). Locusts were fed bran and leaves of *Bromus catharticus* grass or sorghum grown at a research plot at the National Institute of Agro-biological Sciences at Ohwashi in Tsukuba. They laid several egg pods over a month's interval. These eggs were incubated at 20 °C for a month before being chilled at 5 °C for 3 months to terminate embryonic diapause (Tanaka 1992). Eggs chilled were then incubated at 30 °C for hatching. Approximately 100 nymphs, derived from 3 to 4 egg pods laid by the same female, were reared in a large cage (42 × 42 × 24 cm) and 34 family lines were thus obtained. More than 15 egg pods were collected from each family line and exposed to a series of temperature regimes to terminate diapause as described above. Green hatchlings appeared within two such family lines when a total of 12 egg pods were incubated at 30 °C for hatching. Unfortunately, the incidence of green hatchlings for each egg pod was not recorded accurately, but they appeared together with normal-colored individuals from the same egg pods. Green and normal-colored nymphs were reared together in a large cage under the same conditions as above.

## Results

As mentioned above, 2 out of 34 family lines produced green hatchlings in the 2<sup>nd</sup> generation. Fig. 1a shows newly hatched nymphs obtained from one egg pod. Two types of hatchlings could be recognized: one type had a light-brown body color with black eyes shortly after hatching (Fig. 1d): the body turned dark brown several hours later (Fig. 1e); the other type exhibited a yellow-green body color with reddish eyes in the beginning (Fig. 1b) and developed a darker body color later (Fig. 1c). (The latter type will be called a 'green morph'.) After ecdysis to the 2<sup>nd</sup> instar, green morphs assumed a light body color (Fig. 1f) compared with normal locusts (Fig. 1g).

Mortality was considerably higher among green morphs and only 7 of the 22 individuals reared (32%) reached the last nymphal instar, whereas more than 70% developed to the same stage among normal-colored locusts reared in the same cage (total N = 25). Last instar nymphs of the green morph were slightly greenish with light-colored compound eyes (Fig. 1h).

Two females and two males of this morph emerged as adults and were transferred to another cage. Of these, one female and one male died a few days after adult emergence. The surviving two individuals still remained whitish two weeks after adult emergence (Fig. 1j). This 'pair' started mating in two weeks, but the female died without producing any eggs. The male survivor was then kept with two normal female adults (Fig. 1k) to obtain eggs. Mating was witnessed a few times, but no viable eggs were obtained from these females.




Fig. 1. Photographs showing green and normal morphs of *L. migratoria*. a. Hatchlings of green and normal morph. b. Less than hour-old green hatchling. c. Green hatchling a few hours old. d. Less than hour-old normal hatchling. e. Normal hatchling a few hours old. f. 2<sup>nd</sup> instar green nymph. g. 2<sup>nd</sup> instar normal nymph. h. Last (5<sup>th</sup>) instar green nymph. i. Last (5<sup>th</sup>) instar normal nymph displaying a typical gregarious color. j. Green morph adult. k. Normal morph adult. Horizontal bars indicate 1cm. For color version, see Plate II.

## Discussion

The green morphs observed in the present study exhibited body coloration distinctly different from that of their normal-colored counterparts reared in the same cage. Green hatchlings are common among locusts and grasshoppers (Uvarov 1977). In *L. migratoria*, hatchling body color shows some variation, depending on the parental phase or crowding conditions (Hunter-Jones 1958), but green-colored hatchlings are not known. *L. migratoria* albinism has been reported from different laboratories (Faure 1932, Verdier 1965, Nolte 1969, Hasegawa & Tanaka 1996). The albino strain from Okinawa is controlled by a single recessive Mendelian unit and deficient in the dark-color inducing hormone (Tanaka 1993, Tanaka & Pener 1994) now known as [His<sup>7</sup>]-corazonin (Tawfik *et al.* 1999). Hatchlings of this albino mutant are, however, indistinguishable from normal hatchlings by either body color or eye color: they have

a fawn body color and black eyes (Tanaka S., unpub. obs.). In the desert locust *Schistocerca gregaria* Forskål, an albino strain controlled by a simple recessive Mendelian unit is also known (Hunter-Jones 1957). Yerushalmi *et al.* (2000) obtained a "dark-adult" mutant in this same locust that is also recessive to the normal phenotype. Nymphs of this "dark-adult" strain look the same as normal hoppers, but develop a black pigment in the body and wings after adult emergence.

The green morph of *L. migratoria* suffered from a high rate of mortality in the laboratory compared with normal-colored individuals from the same egg pods. Although no genetic studies have been performed, it seems likely that the green morphs carried a semilethal gene. As mentioned earlier, the incidence of green hatchlings was not recorded accurately; however, they always appeared together with normal phenotypes from the same egg pods. These observations, together with the fact that they did not appear until the 2<sup>nd</sup> genera-



tion in inbred family lines, might suggest that the green-morph phenotype was controlled by a recessive, possibly semilethal gene.

Recently we obtained a short-winged morph of *L. migratoria* in a similar way (Tanaka & Nishide 2012): several short-winged individuals appeared in the 2<sup>nd</sup> generation of inbred family lines originating from Tsushima Island, Japan. The short-winged phenotype is controlled by a single recessive Mendelian unit. Although the possibility that these unexpected morphs have occurred by mutation in the laboratory cannot be excluded, inbreeding wild-caught insects in the laboratory may be an interesting way of finding recessive phenotypes that are normally unexpressed in the field.

### Acknowledgements

The authors thank Ms N. Totsuka, Ms M. Higuchi and Ms Y. Yokota for assistance with rearing locusts. This study was supported in part by a Kakenhi Grant (no. 23380038). The grass used in this experiment was raised by the Field Management Section of the National Institute of Agro-biological Sciences at Ohwashi. An anonymous reviewer improved the manuscript.

### References

- Faure J.C. 1932. The phases of locusts in South Africa. *Bulletin of Entomological Research* 23: 293-405.
- Hakomori T., Tanaka S. 1992. Genetic control of diapause and other developmental traits in Japanese strains of the migratory locust, *Locusta migratoria*: univoltine vs bivoltine. *Japanese Journal of Entomology* 60: 319-328.
- Hasegawa E., Tanaka S. 1996. Sexual maturation in *Locusta migratoria* females: laboratory vs field conditions. *Applied Entomology and Zoology* 31: 279-290.
- Hunter-Jones P. 1957. An albino strain of the desert locust. *Nature* 180: 236-237.
- Hunter-Jones P. 1958. Laboratory studies on the inheritance of phase characters in locusts. *Anti-Locust Bulletin* 29: 1-32.
- Nolte D.J. 1969. Chiasma induction and tyrosine metabolism in locusts. *Chromosoma (Berlin)* 26: 287-297.
- Pener M.P. 1991. Locust phase polyphenism and its endocrine relations. *Advances in Insect Physiology* 23: 1-79.
- Pener M.P., Yerushalmi, Y. 1998. The physiology of locust phase polymorphism: an update. *Journal of Insect Physiology* 44: 365-377.
- Pener M.P., Simpson J.S. 2009. Locust phase polyphenism: an update. *Advances in Insect Physiology* 36: 1-272.
- Tanaka S. 1992. The significance of embryonic diapause in a Japanese strain of the migratory locust, *Locusta migratoria* (Orthoptera: Acrididae). *Japanese Journal of Entomology* 60: 503-520.
- Tanaka S. 1993. Hormonal deficiency causing albinism in *Locusta migratoria*. *Zoological Science* 10: 467-471.
- Tanaka S., Nishide Y. 2012. First record of the occurrence and genetics of a short-winged morph in the migratory locust, *Locusta migratoria* (Orthoptera: Acrididae). *Journal of Orthoptera Research* 21: 169-174.
- Tanaka S., Pener M.P. 1994. A neuropeptide controlling the dark pigmentation in color polymorphism of the migratory locust, *Locusta migratoria*. *Journal of Insect Physiology* 40: 997-1005.
- Tawfik A.I., Tanaka S., De Loof A., Schoofs L., Baggerman G., Waelkens E., Derua R., Milner Y., Yerushalmi Y., Pener M.P. 1999. Identification of the gregarization-associated dark-pigmentotropin in locusts through an albino mutant. *Proceedings National Academy of Science USA* 96: 7083-7087.
- Uvarov B.P. 1966. *Grasshoppers and Locusts*, Vol. 1. Cambridge: Cambridge University Press. 481 pp.
- Uvarov B.P. 1977. *Grasshoppers and Locusts*, Vol. 2. Centre for Overseas Pest Research, London. 613 pp.
- Verdier M. 1965. Mutation albinos de *Locusta migratoria*. 1. Origine et description (C.S). *Bulletin Société Zoologique France* 90: 493-501.
- Yerushalmi Y., Abu-Hilal H., Pener M.P. 2000. A "dark-adult" mutation of *Schistocerca gregaria* (Forsk.). *Journal of Orthoptera Research* 9: 41-43.